Projektbereich B6

Discussion Paper No. B-361

Why Imitate, and if so, How? A Bounded Rational Approach to Multi-Armed Bandits[1]

Karl H. Schlag[2]

Febuary, 1996

[1]The author wishes to thank Dirk Bergemann, Jonas Björnerstedt, Georg Nöldeke, Larry Samuelson, Avner Shaked, a referee and an associate editor for helpful comments. Financial support from the Deutsche Forschungsgemeinschaft, Sonderforschungsbereich 303 at the University of Bonn is gratefully acknowledged.

[2]Abt. Wirtschaftstheorie III, Department of Economics, University of Bonn, Adenauerallee 24-26, 53113 Bonn, Germany.

Abstract

We consider the situation in which individuals in a finite population must repeatedly choose an action yielding an uncertain payoff. Between choices, each individual may observe the performance of one other individual. We search for rules of behavior with limited memory that increase expected payoffs for any underlying payoff distribution. It is shown that the rule that outperforms all other rules with this property is the one that specifies imitation of the action of an individual that performed better with a probability proportional to how much better she performed. When each individual uses this best rule, the aggregate population behavior can be approximated by the replicator dynamic.

JEL Classification Number: C72, C79

Keywords: social learning, bounded rationality, imitation, multi-armed bandit, random matching, payoff increasing, replicator dynamic.

1 Introduction

Imitation is the act of copying or mimicking the action of others, a commonly observed behavior in human decision making. This is perhaps the reason why lately, with the growing popularity of Evolutionary Game Theory, many models of social learning have emerged in which individuals select their future actions by imitating others (e.g., Banerjee [1]; Björnerstedt and Weibull [3]; Cabrales [5]; Ellison and Fudenberg [7]; Gale et al. [8]; Hofbauer [9]; Rogers [12]). We propose to answer why individuals should follow an imitative behavior in a social learning environment. Especially, we want to point out that the answer is directly related to the answer to how imitation should take place. Our approach follows the traditional line of decision and game theory in that we allow individuals to determine their behavior before we derive implications of individual behavior on the evolution of the society. At the same time we depart from traditional decision theory (Savage [16]) and suggest two alternative approaches to selecting individual behavior, a bounded rational approach and a population-oriented approach. What is interesting is that either approach leads to the same unique prescription of how to choose future actions:

- follow an imitative behavior, i.e., only change actions through imitating others,
- never imitate an individual that performed worse than oneself, and
- imitate an individual that performed better with a probability that is proportional to how much better this individual performed.

This result is not only interesting for its own sake but also for its implications on the global adjustment of a large population in which each individual follows the individually most preferred rule. In this regard, it turns out that the global process governing the change of actions in the population can be approximated by the well known replicator dynamic (Taylor [20]). In this dynamic, the growth rate of an action is equal to its relative payoff measured with respect to the average payoff in the population.

This paper generalizes an earlier work by the author (Schlag [17]) to a multi-armed bandit setting. Individuals must repeatedly choose an action from a finite set of actions A. Choosing an action yields an uncertain payoff. Payoffs are realized independently, their distribution has finite support and belongs to a bounded interval $[\alpha, \omega]$. This basic setting of multi-armed bandits has a wide range of applications to economics and behavioral sciences; choosing an action can be a synonym for the choice of a technology, of an organizational structure or of a restaurant, it may be the setting of a price (Ellison and Fudenberg [7]; Schmalensee [19]). In our model, individuals belong to a finite population and are able to learn from others. After each payoff realization each individual is able to observe (or sample) the present performance of another individual; she observes the action chosen and the payoff received by the sampled individual. Sampling occurs according to some exogenously given distribution. However, we do require that this procedure of sampling is symmetric, i.e., that individual 'c' samples individual 'd' with the same probability as individual 'd' samples individual 'c'. Random sampling will play a special role regarding aggregate population behavior.

The action chosen by an individual in the next round as a function of the information that she has acquired in the past will be called a *behavioral rule*. We want to focus on simple behavioral rules and thus limit the amount of information available to the individual about previous occurrences as follows. We assume that an individual forgets all about actions and observations from previous rounds. Hence, the rule determining play in the next round is a function of the payoffs achieved and actions taken both by oneself and the sampled individual in the present round. Moreover, we assume that an individual faces many multi-armed bandits that she can not distinguish a priori except for the fact that they all have the same set of possible actions. Hence, an individual must follow the same behavior (use the same behavioral rule) whenever she is faced with a multi-armed bandit with the same set of actions. Especially we have implicitly ruled out an individual that has enough memory capacity to be able to choose future actions by updating some prior. Similarly we have ruled out complex strategies as used in the classical multi-armed bandit problem (e.g., see Rothschild [13]). In the following consider the set of actions A and the interval $[\alpha, \omega]$ containing the realized payoffs fixed. As mentioned above, we follow two alternative approaches for selecting a behavioral rule for our setting.[3]

The first approach (formalized in Section 4.1) is from the standpoint of a boundedly rational individual. The individual enters a finite population of individuals who each face a multi-armed bandit. All multi-armed bandits yield the same payoff distributions. Entry occurs by replacing a randomly selected individual and adapting the action this individual chose in the last round. As mentioned above, the individual must use the same behavioral rule when facing each multi-armed bandit. The following assumptions are made. The individual measures her success in a given decision situation by the expected increase (or decrease) in payoffs her present choice yields in comparison to her previous choice. The individual wants to perform well in each situation in contrast to performing badly in some and very well in others. Hence, we aim to investigate if there is a rule in our setting that always leads to an increase in expected payoffs. At this point we add an additional bounded rational assumption, namely, we assume that the entering individual treats each round as if she had just entered. This means that the individual does not anticipate how her decisions affect future distributions of actions played in the population. A rule that (weakly) increases expected payoffs in any decision situation, i.e., in any state regardless of the underlying payoff distributions associated to the various actions, under the above assumption of ignorance of previous play, will be called *improving*. If an improving rule induces constant expected payoffs in all decision situations then the rule will be called *stationary*.

Our second approach to selecting behavioral rules (formalized in Section 4.2) takes a population-oriented perspective. In this scenario we will search

[3]Unless otherwise stated, we ignore the issue of which action to choose when the individual has no prior experience.

for a rule that, when played by each individual in a finite population, is expected to (weakly) increase average payoffs in the population over time in any decision situation. Such a rule will be called *payoff increasing*. For large (i.e., infinite) populations, this means that we search for a rule that induces a better reply dynamic (Hofbauer [9]) in all decision situations when the population is made up of replicas of the individual using this rule. In a model that includes evolutionary pressure on behavioral rules it is plausible that only payoff increasing rules can survive the entry of an alternative rule in all environments. Björnerstedt and Schlag [2] confirm this intuition, a result explained further in Appendix B.

The two approaches above have in common that a certain property must hold in all decision situations. A fundamental connection between these two approaches is that a rule is improving in the boundedly rational model if and only if it is payoff increasing in the population model. Hence, from the standpoint of either selection approach our initial aim will be to characterize the improving rules. Trivial examples of stationary improving rules are easily given, e.g., the self-explanatory rules 'never switch' and 'always switch'. We note that an improving rule is stationary if and only if it always induces constant average payoffs in the population. For this reason, we will focus special attention on the class of non-stationary improving rules.

One important aspect of our first lemma is that improving rules are imitating, i.e., that switching actions occurs by imitating the sampled individual. From this, the main theorem (Theorem 1) characterizes the set of improving rules. The criterion for an imitating rule to be improving is that the difference in the probabilities of switching when two individuals sample each other is *proportional* to the difference in the payoffs each individual achieved. The reason for this strong condition on the probabilities of switching relies on the linear structure of taking expectations. An intuitive and hence popular behavioral rule is the rule we call *'imitate if better'* (used e.g. by Ellison and Fudenberg [7] and Malawski [10]). This rule prescribes to imitate the action of the observed individual whenever she achieves a higher payoff. Our analysis reveals that this rule is not improving. Another consequence of our main theorem is that a bound on the set of feasible payoffs (in our model, payoffs are contained in $[\alpha, \omega]$) is necessary to obtain non stationary improving rules.

Our classification of the improving rules reveals a clear structure among the improving rules. This leads in both selection approaches to a natural criterion for selecting a best rule among the improving rules. In fact, in either selection approach the same behavioral rule is selected. This rule is a specific rule from the class of rules we call *proportional imitation rules* that were described at the beginning of this introduction. Especially, we obtain a unique prescription of how to choose actions in our model.

Next we investigate implications for the population adjustment process under random sampling when each individual is using our prescribed best behavioral rule. We show that the process governing the adaptation of actions in a large population can be approximated in the short run by a discrete version of the replicator dynamic (Taylor [20]) applied to this setting. Especially, with probability arbitrarily close to one, most individuals will be playing after a finite number of rounds an action that is best (measured in terms of expected payoffs) among those initially present.

In a further section we consider a two population random matching scenario. All the previous analysis and results are shown to apply to this generalized setting. Especially, individuals prefer to use a specific proportional imitation rule. Moreover, in an infinite population under random sampling in which each individual uses the optimal individual behavior, the adjustment is described by a discrete version of the replicator dynamic (Taylor [20]).

The rest of the paper is organized as follows. In Section two the basic payoff realization and sampling scenario is introduced. The feasible behavioral rules for using in this scenario are presented in Section three. Section four contains two alternative approaches to selecting a behavioral rule, each leading to the condition of improving. In Section five we present a first lemma concerning improving rules. In Section six this lemma is applied to show that 'imitate if better' is not improving. Section seven contains the central theorem of this paper, a complete characterization of an improving rule. In Section eight we select among the improving rules. Section nine contains the implications this has for the population dynamic. In Section ten we consider an alternative two population matching scenario. Section twelve contains a discussion. Appendix A contains some the proof of the main theorem. In appendix B some in sights to the evolutionary model of Björnerstedt and Schlag [2] are given.

2 The Payoff Realization and Sampling Scenario

Consider the following dynamic process of choosing actions, sampling and updating. Let P denote the underlying probability measure. Let W be a finite population (or set) of N individuals, $N \geq 2$. In a sequence of rounds, each individual must choose an action from a finite set of actions $A = \{1, .., n\}$, $n \geq 2$. Choosing the action i yields an uncertain payoff drawn from a probability distribution P_i with finite support in $[\alpha, \omega]$ where α and ω, $\alpha < \omega$, are exogenous parameters. Payoffs are realized independently of all other events. Let π_i denote the expected payoff generated by choosing action i, i.e., $\pi_i = \sum_{x \in [\alpha,\omega]} x P_i(x)$, $i \in A$. Then the tuple $\left\langle A, (P_i)_{i \in A} \right\rangle$ constitutes a *multi-armed bandit* or a game against nature. The set of all multi-armed bandits with action set A yielding payoffs in $[\alpha, \omega]$ will be denoted by $\mathcal{G}(A, [\alpha, \omega])$.

A state $s \in A^W$ of the population in a given round t is the description of the action that each individual is choosing in round t. Let $m_i = m_i(s)$ denote the number of individuals choosing the action i in state s, i.e., $m_i = |\{c \in W : s(c) = i\}|$ $(i \in A)$. Let $\Delta(A)$ be the set of probability distributions on A. For a given state s let $p \in \Delta(A)$ denote the probability distribution that is associated with randomly selecting an individual and observing the action she has chosen for this round, i.e., $p_i = m_i/N$. The set of all such probability distributions will be denoted by $\Delta^N(A)$, i.e., $p \in \Delta^N(A)$ and $i \in A$ implies $N \cdot p_i \in \mathbb{N}$. Given this notation, the average expected payoff of the population in state s, $\bar{\pi}(s)$, is given by $\bar{\pi}(s) = \sum p_i \pi_i$.

Between the rounds of payoff realization, each individual meets (or samples) another individual from the population and receives the following information. When individual 'c' samples individual 'd' $(c, d \in W)$, then individual 'c' observes the action 'd' used and the payoff 'd' achieved in the last round without observing the identity of 'd'. For each individual 'c' this sampling occurs according to some exogenously given probability distribution $z_c \in \Delta(W \setminus \{c\})$ where $z_c(d)$ is the probability that individual 'c' samples individual 'd'. z_c is called a *sampling rule* for individual 'c'. The assignment of a sampling rule z_c to each individual c in the population will be called a sampling procedure. Formally, $z = (z_c)_{c \in W}$ is called a *sampling procedure* if $z_c \in \Delta(W \setminus \{c\})$ for all $c \in W$. A sampling procedure z will be called *symmetric* if for any $c, d \in W$ the probability of 'c' sampling 'd' (this event denoted by $c \rightsquigarrow d$) is the same as vice versa, i.e., $P(c \rightsquigarrow d) = P(d \rightsquigarrow c)$. In the following we will restrict attention to symmetric sampling procedures.

The above conditions restrict the variety of individual sampling procedures without specifying explicitly how the sampling rules of different individuals relate to each other. A scenario in which each individual is sampled at most once (due to time constraints) is equally feasible as one in which an individual can be sampled a multiple number of times. The sampling could be such that individuals sample independently, i.e., $P(c \rightsquigarrow d | d \rightsquigarrow c) = P(c \rightsquigarrow d)$. Similarly, we allow for a model in which individuals sample each other. In this case $c \rightsquigarrow d$ is the same event as $d \rightsquigarrow c$ for each $c, d \in W$, especially, $P(c \rightsquigarrow d | d \rightsquigarrow c) = 1$ and the sampling rule is symmetric. For example, a symmetric sampling procedure results from the following assumptions on individual sampling behavior. Individuals are located on a circle. Each individual randomly samples with equal probability among her $2m$ closest neighbors (m to the left, m to the right, $m < N/2$). Special attention will focus on the symmetric sampling procedures referred to as *random sampling*. Here each individual randomly samples an individual (with equal probability) from the population (except for herself), i.e., $P(c \rightsquigarrow d) = \frac{1}{N-1}$ for $c, d \in W$, $c \neq d$.

3 Behavioral Rules

Let A, $[\alpha, \omega]$ and N be fixed throughout the rest of the paper. The description of how an individual in our model chooses her next action whenever she faces a multi-armed bandit with action set A that realizes payoffs in $[\alpha, \omega]$ is summarized by a *behavioral rule*. Especially, the individual may not change her rule over time. We allow for an individual to use a randomizing device that generates independent events when they determine which action to choose in the next round. Following the assumptions on individual information and memory made in the introduction, a *behavioral rule* F is characterized by a function $F : A \times [\alpha, \omega] \times A \times [\alpha, \omega] \to \Delta(A)$ where $F(i, x, j, y)_k$ is the probability of choosing action k in the next round after previously choosing action i, receiving payoff x and sampling an individual who chose action j and received payoff y. For simplicity we will disregard the issue of which action to choose in the first round.

One of the simplest behavioral rules is the rule '*never switch*', formally defined by the behavioral rule F that satisfies $F(i, x, j, y)_i = 1$ for $i, j \in A$ and $x, y \in [\alpha, \omega]$. An opposite behavior is exhibited by the self explanatory rule '*always switch*'. A more plausible rule seems to be to act according to '*imitate if better*' (Ellison and Fudenberg [7]; Malawski [10]), i.e., use the rule F given by $F(i, x, j, y)_j = 1$ if $y > x$ and $F(i, x, j, y)_i = 1$ if $y \leq x$. The three rules described above belong to the class of behavioral rules that are based on imitation, i.e., either the individual does not change actions or she switches to the action used by the individual she sampled. More generally, we call a behavioral rule F *imitating* if $F(i, x, j, y)_k = 0$ when $k \notin \{i, j\}$ $(x, y \in [\alpha, \omega])$. The class of imitating rules, referred to as proportional imitation rules, described at the beginning of the introduction will play an important role in our analysis. A behavioral rule F is called a *proportional imitation rule* if F is imitating and there exists σ with $0 < \sigma \leq \frac{1}{\omega - \alpha}$ such that $F(i, x, j, y)_j = 0$ if $y \leq x$ and $F(i, x, j, y)_j = \sigma(y - x)$ if $y > x$, $i \neq j$ and $x, y \in [\alpha, \omega]$. The constant σ associated to a given proportional imitation rule will be referred to as the rate.[4]

[4] Switching behavior as displayed by proportional imitation rules appears in a paper by Cabrales [5] where it is justified through uniformly distibuted costs for switching actions.